

Testing supersymmetric models with direct detection of dark matter

Paweł Szczerbiak

University of Warsaw

Astrofizyka Cząstek w Polsce 2017


September 21, 2017, Kraków, UJ

In collaboration with M. Badziak, M. Olechowski



Based on: 1: Phys. Lett. B **770** (2017) 226-235

2: JHEP **1707** (2017) 050




Motivation: direct detection of dark matter



Motivation: direct detection of dark matter



Motivation: direct detection of dark matter



Models and basic constraints

Models:

- ▶ Well-tempered bino-Higgsino in MSSM
- ▶ Singlino-Higgsino in NMSSM

$$N_{12} \approx 0$$

$$N_{11}, N_{12} \approx 0$$

Models and basic constraints

Models:

- ▶ Well-tempered bino-Higgsino in MSSM
- ▶ Singlino-Higgsino in NMSSM

$$N_{12} \approx 0$$

$$N_{11}, N_{12} \approx 0$$

Constraints:

- ▶ $\Omega h^2 \approx 0.12$

- ▶ $\sigma_{\text{SI}} \sim \left(\frac{\alpha_{\chi\chi h} \alpha_{hNN}}{2m_h^2} + \frac{\alpha_{\chi\chi H} \alpha_{HNN}}{2m_H^2} + \dots \right)^2 \lesssim \sigma_{\text{SI}}^{\text{LUX}}$

- ▶ $\sigma_{\text{SD}} \sim (N_{13}^2 - N_{14}^2)^2 \lesssim \sigma_{\text{SD}}^{\text{LUX}}$

Models and basic constraints

Models:

- ▶ Well-tempered bino-Higgsino in MSSM $N_{12} \approx 0$
- ▶ Singlino-Higgsino in NMSSM $N_{11}, N_{12} \approx 0$

Constraints:

- ▶ $\Omega h^2 \approx 0.12$
- ▶ $\sigma_{\text{SI}} \sim \left(\frac{\alpha_{\chi\chi h} \alpha_{hNN}}{2m_h^2} + \frac{\alpha_{\chi\chi H} \alpha_{HNN}}{2m_H^2} + \dots \right)^2 \lesssim \sigma_{\text{SI}}^{\text{LUX}}$
- ▶ $\sigma_{\text{SD}} \sim (N_{13}^2 - N_{14}^2)^2 \lesssim \sigma_{\text{SD}}^{\text{LUX}}$

$$N_{13}^2 - N_{14}^2 = \frac{\left[1 - (m_\chi/\mu)^2 \right] (1 - N_{11(15)}^2) \cos 2\beta}{1 + (m_\chi/\mu)^2 + 2(m_\chi/\mu) \sin 2\beta}$$

Models and basic constraints

Models:

- ▶ Well-tempered bino-Higgsino in MSSM $N_{12} \approx 0$
- ▶ Singlino-Higgsino in NMSSM $N_{11}, N_{12} \approx 0$

Constraints:

- ▶ $\Omega h^2 \approx 0.12$
- ▶ $\sigma_{\text{SI}} \sim \left(\frac{\alpha_{\chi\chi h} \alpha_{hNN}}{2m_h^2} + \frac{\alpha_{\chi\chi H} \alpha_{HNN}}{2m_H^2} + \dots \right)^2 \lesssim \sigma_{\text{SI}}^{\text{LUX}}$
- ▶ $\sigma_{\text{SD}} \sim (N_{13}^2 - N_{14}^2)^2 \lesssim \sigma_{\text{SD}}^{\text{LUX}}$

$$N_{13}^2 - N_{14}^2 = \frac{\left[1 - (m_\chi/\mu)^2\right] (1 - N_{11(15)}^2) \cos 2\beta}{1 + (m_\chi/\mu)^2 + 2(m_\chi/\mu) \sin 2\beta}$$



SI blind spots possible

SD blind spots only for $\tan \beta = 1$ or pure states

Well-tempered bino-Higgsino in MSSM – heavy H

$$\alpha_{h\chi\chi}\alpha_{hNN} \sim \frac{N_{11}^2}{\mu} \frac{m_\chi/\mu + \sin 2\beta}{1 - (m_\chi/\mu)^2} \quad \alpha_{H\chi\chi}\alpha_{HNN} \sim 0$$



$$\sigma_{\text{SI}} = 0 \quad \text{for} \quad \frac{m_\chi}{\mu} + \sin 2\beta \approx 0$$



Well-tempered bino-Higgsino in MSSM – light H

$$\alpha_{h\chi\chi}\alpha_{hNN} \sim \frac{N_{11}^2}{\mu} \frac{m_\chi/\mu + \sin 2\beta}{1 - (m_\chi/\mu)^2} \quad \alpha_{H\chi\chi}\alpha_{HNN} \sim \frac{N_{11}^2}{\mu} \frac{\text{tg } \beta}{1 - (m_\chi/\mu)^2}$$

$$\sigma_{\text{SI}} = 0 \quad \text{for} \quad \frac{m_\chi}{\mu} + \sin 2\beta \approx -\frac{m_h^2}{m_H^2} \frac{\text{tg } \beta}{2}$$



Singlino-Higgsino in NMSSM

$$\begin{aligned} W &= \lambda S H_u H_d + \xi_F S + \frac{1}{2} \mu' S^2 + \frac{1}{3} \kappa S^3 \\ -\mathcal{L}_{\text{soft}} &\supset m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 \\ &+ A_\lambda \lambda H_u H_d S + \frac{1}{3} A_\kappa \kappa S^3 + m_3^2 H_u H_d + \frac{1}{2} m_\xi^2 S^2 + \xi_S S + h.c. \end{aligned}$$

Singlino-Higgsino in NMSSM

$$\begin{aligned} W &= \lambda S H_u H_d + \xi_F S + \frac{1}{2} \mu' S^2 + \frac{1}{3} \kappa S^3 \\ -\mathcal{L}_{\text{soft}} &\supset m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 \\ &+ A_\lambda \lambda H_u H_d S + \frac{1}{3} A_\kappa \kappa S^3 + m_3^2 H_u H_d + \frac{1}{2} m_\xi^2 S^2 + \xi_S S + h.c. \end{aligned}$$

Considered cases:

1. Only h exchange
 - ▶ no mixing among scalars
 - ▶ with scalar mixing
2. h and H exchange
 - ▶ no mixing with s
 - ▶ mixing with s , $m_s \gg m_h$
3. h and s exchange
 - ▶ leading effect from H

Singlino-Higgsino in NMSSM

$$\begin{aligned} W &= \lambda S H_u H_d + \xi_F S + \frac{1}{2} \mu' S^2 + \frac{1}{3} \kappa S^3 \\ -\mathcal{L}_{\text{soft}} &\supset m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 \\ &+ A_\lambda \lambda H_u H_d S + \frac{1}{3} A_\kappa \kappa S^3 + m_3^2 H_u H_d + \frac{1}{2} m_\xi'^2 S^2 + \xi_S S + h.c. \end{aligned}$$

Considered cases:

1. Only h exchange

- ▶ no mixing among scalars $\frac{m_\chi}{\mu} - \sin 2\beta = 0$ (MSSM-like)
- ▶ with scalar mixing

2. h and H exchange

- ▶ no mixing with s $\frac{m_\chi}{\mu} - \sin 2\beta = \frac{\tan \beta}{2} \left(\frac{m_h}{m_H} \right)^2$ (MSSM-like)
- ▶ mixing with s , $m_s \gg m_h$

3. h and s exchange

- ▶ leading effect from H

Singlino-Higgsino in NMSSM

$$\begin{aligned} W &= \lambda S H_u H_d + \xi_F S + \frac{1}{2} \mu' S^2 + \frac{1}{3} \kappa S^3 \\ -\mathcal{L}_{\text{soft}} &\supset m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 \\ &+ A_\lambda \lambda H_u H_d S + \frac{1}{3} A_\kappa \kappa S^3 + m_3^2 H_u H_d + \frac{1}{2} m_\xi^2 S^2 + \xi_S S + h.c. \end{aligned}$$

Considered cases:

1. Only h exchange in this talk

- ▶ no mixing among scalars $\frac{m_\chi}{\mu} - \sin 2\beta = 0$ (MSSM-like)
- ▶ with scalar mixing

2. h and H exchange

- ▶ no mixing with s $\frac{m_\chi}{\mu} - \sin 2\beta = \frac{\tan \beta}{2} \left(\frac{m_h}{m_H} \right)^2$ (MSSM-like)
- ▶ mixing with s , $m_s \gg m_h$

3. h and s exchange in this talk

- ▶ leading effect from H

General NMSSM – only h exchange

Two generic mechanisms:

- ▶ resonance with Z^0 boson ($m_\chi \sim 45$ GeV):

$$\Omega h^2 \approx 0.1 \left(\frac{0.3}{N_{13}^2 - N_{14}^2} \right)^2 \frac{m_Z^2}{4m_\chi^2} \left[\left(\frac{4m_\chi^2}{m_Z^2} - 1 + \frac{\bar{v}^2}{4} \right)^2 + \frac{\Gamma_Z^2}{m_Z^2} \right]$$

- ▶ annihilation into $t\bar{t}$ ($m_\chi \gtrsim 170$ GeV):

$$\Omega h^2 \approx 0.1 \left(\frac{0.05}{N_{13}^2 - N_{14}^2} \right)^2 \left[\sqrt{1 - \frac{m_t^2}{m_\chi^2}} + \frac{3}{4} \frac{1}{x_f} \left(1 - \frac{m_t^2}{2m_\chi^2} \right) \frac{1}{\sqrt{1 - \frac{m_t^2}{m_\chi^2}}} \right]^{-1/2}$$

General NMSSM – only h exchange

Two generic mechanisms:

- ▶ resonance with Z^0 boson ($m_\chi \sim 45$ GeV):

$$\Omega h^2 \approx 0.1 \left(\frac{0.3}{N_{13}^2 - N_{14}^2} \right)^2 \frac{m_Z^2}{4m_\chi^2} \left[\left(\frac{4m_\chi^2}{m_Z^2} - 1 + \frac{\bar{v}^2}{4} \right)^2 + \frac{\Gamma_Z^2}{m_Z^2} \right]$$


- ▶ annihilation into $t\bar{t}$ ($m_\chi \gtrsim 170$ GeV):

$$\Omega h^2 \approx 0.1 \left(\frac{0.05}{N_{13}^2 - N_{14}^2} \right)^2 \left[\sqrt{1 - \frac{m_t^2}{m_\chi^2}} + \frac{3}{4} \frac{1}{x_f} \left(1 - \frac{m_t^2}{2m_\chi^2} \right) \frac{1}{\sqrt{1 - \frac{m_t^2}{m_\chi^2}}} \right]^{-1/2}$$


$$\sigma_{\text{SD}} \sim (N_{13}^2 - N_{14}^2)^2$$

General NMSSM – only h exchange

Resonance with Z^0




Annihilation into $t\bar{t}$




General NMSSM – only h exchange

Resonance with Z^0



Annihilation into $t\bar{t}$



Blind spot conditions:

► No mixing among scalars: $\frac{m_\chi}{\mu} - \sin 2\beta \approx 0$

► With scalar mixing: $\frac{m_\chi}{\mu} - \sin 2\beta \approx \gamma \frac{\kappa}{\lambda} \left(\frac{\mu}{\lambda v} \right) \left(1 - \left(\frac{m_\chi}{\mu} \right)^2 \right)$

Where: $\gamma \equiv \frac{\tilde{S}_{h\hat{s}}}{\tilde{S}_{h\hat{h}}} \sim \sqrt{|\Delta_{\text{mix}}|}$

General NMSSM – h and s exchange

- Let us introduce:

$$\mathcal{A}_s \equiv \frac{\alpha_{sNN}}{\alpha_{hNN}} \frac{\tilde{S}_{s\hat{s}}}{\tilde{S}_{h\hat{h}}} \left(\frac{m_h}{m_s} \right)^2 \quad \gamma \equiv \frac{\tilde{S}_{h\hat{s}}}{\tilde{S}_{h\hat{h}}} \sim \sqrt{|\Delta_{\text{mix}}|}$$

General NMSSM – h and s exchange

- ▶ Let us introduce:

$$\mathcal{A}_s \equiv \frac{\alpha_{sNN}}{\alpha_{hNN}} \frac{\tilde{S}_{s\hat{s}}}{\tilde{S}_{h\hat{h}}} \left(\frac{m_h}{m_s} \right)^2 \quad \gamma \equiv \frac{\tilde{S}_{h\hat{s}}}{\tilde{S}_{h\hat{h}}} \sim \sqrt{|\Delta_{\text{mix}}|}$$

- ▶ Blind spot condition:

$$\frac{m_\chi}{\mu} - \sin 2\beta \approx \frac{\gamma + \mathcal{A}_s}{1 - \gamma \mathcal{A}_s} \frac{\kappa}{\lambda} \left(\frac{\mu}{\lambda v} \right) \left(1 - \left(\frac{m_\chi}{\mu} \right)^2 \right)$$

General NMSSM – h and s exchange

- ▶ Let us introduce:

$$\mathcal{A}_s \equiv \frac{\alpha_{sNN}}{\alpha_{hNN}} \frac{\tilde{S}_{s\hat{s}}}{\tilde{S}_{h\hat{h}}} \left(\frac{m_h}{m_s} \right)^2 \quad \gamma \equiv \frac{\tilde{S}_{h\hat{s}}}{\tilde{S}_{h\hat{h}}} \sim \sqrt{|\Delta_{\text{mix}}|}$$

- ▶ Blind spot condition:

$$\frac{m_\chi}{\mu} - \sin 2\beta \approx \frac{\gamma + \mathcal{A}_s}{1 - \gamma \mathcal{A}_s} \frac{\kappa}{\lambda} \left(\frac{\mu}{\lambda v} \right) \left(1 - \left(\frac{m_\chi}{\mu} \right)^2 \right)$$

- ▶ Conclusions:



- ▶ because $m_s < m_h$, the RHS can be one order of magnitude larger as compared to the case of only h exchange
- ▶ in general NMSSM we can have $\Omega h^2 \approx 0.12$ and other experimental bounds fulfilled even for $\Delta_{\text{mix}} \sim 4 \text{ GeV}$, where $\gamma \sim \sqrt{\Delta_{\text{mix}}}$, $m_h = \hat{M}_{hh} + \Delta_{\text{mix}}$.

$\mathbb{Z}_3$ -NMSSM – only h exchange (heavy singlet)

$$\frac{m_\chi}{\mu} - \sin 2\beta \approx \gamma \frac{\kappa}{\lambda} \left(\frac{\mu}{\lambda v} \right) \left(1 - \left(\frac{m_\chi}{\mu} \right)^2 \right)$$

$$\text{sgn}(m_\chi \mu) = \text{sgn}(\kappa) \quad |\kappa| < \frac{1}{2} \lambda \quad (\text{for singlino-like LSP})$$

$$m_s^2 + \frac{1}{3} m_a^2 \approx m_{\text{LSP}}^2 + \gamma^2 (m_s^2 - m_h^2) \quad \Rightarrow \quad m_{\text{LSP}} > m_s$$





$\mathbb{Z}_3$ -NMSSM – h and s exchange (light singlet)

$$\frac{m_\chi}{\mu} - \sin 2\beta \approx \frac{\gamma + \mathcal{A}_s}{1 - \gamma \mathcal{A}_s} \frac{\kappa}{\lambda} \frac{\mu}{\lambda v_h} \left(1 - \left(\frac{m_\chi}{\mu} \right)^2 \right)$$

$$m_a \approx 2m_{\text{LSP}} \quad \Rightarrow \quad m_s^2 + \frac{1}{3}m_{\text{LSP}}^2 + \gamma^2(m_h^2 - m_s^2) \approx \Delta_{\hat{s}s} + \frac{1}{3}\Delta_{\hat{a}a}$$

$$m_s^2 \approx m_{\text{LSP}}^2 \left[\left(\frac{\lambda \tan \beta}{2\pi} \right)^2 \ln \left(\frac{2M_{\text{SUSY}}}{m_{\text{LSP}} \tan \beta} \right) - \frac{1}{3} \right]$$



Conclusions

- ▶ We derived current constraints and prospects for SD direct detection for SI blind spots for **bino-Higgsino** LSP in **MSSM** and for **singlino-Higgsino** LSP in **NMSSM**.
- ▶ For **bino-Higgsino**, if H is decoupled, current DD limits set a lower bound $m_{\tilde{t}} \gtrsim 25 \text{ TeV}$. For $H \sim 400 \text{ GeV}$ light stops are possible and this scenario will be tested in near future.
- ▶ For **singlino-Higgsino** the allowed parameter space is still large. If $m_H, m_s \gg m_h$ the allowed mass regions are $m_{\text{LSP}} \sim 41 - 46$ and $300 - 800 \text{ GeV}$ and will be almost entirely probed by XENON1T.
- ▶ In $\mathbb{Z}_3$ -**NMSSM** additional annihilation channels and resonances will relax the SD bounds. In particular, $m_{\text{LSP}} \gtrsim 400 \text{ GeV}$ may not be explored by XENON1T.
- ▶ In the above scenarios, **future SD** limits will play a crucial role in probing the parameter space.

Backup slides: Higgs sector

- Convenient basis ($\hat{H} = \mathcal{O}_\beta H$):

$$\begin{pmatrix} \hat{h} \\ \hat{H} \\ \hat{s} \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta & 0 \\ \sin \beta & -\cos \beta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} H_d \\ H_u \\ S \end{pmatrix}$$

- Mass eigenstates:

$$h_i = \tilde{S}_{ij} \hat{H}_j = S_{ij} H_j \quad \Longrightarrow \quad \tilde{S} = S \cdot \mathcal{O}_\beta$$

Explicitly:

$$h_i = \tilde{S}_{h_i \hat{h}} \hat{h} + \tilde{S}_{h_i \hat{H}} \hat{H} + \tilde{S}_{h_i \hat{s}} \hat{s}$$

$$h_i \equiv h, H, s$$

Backup slides: Higgs sector

- Diagonalization ($\Lambda \equiv A_\lambda + \langle \partial_S^2 f \rangle$):

$$\begin{pmatrix} M_{\hat{h}\hat{h}}^2 & M_{\hat{h}\hat{H}}^2 & M_{\hat{h}\hat{s}}^2 \\ M_{\hat{h}\hat{H}}^2 & M_{\hat{H}\hat{H}}^2 & M_{\hat{H}\hat{s}}^2 \\ M_{\hat{h}\hat{s}}^2 & M_{\hat{H}\hat{s}}^2 & M_{\hat{s}\hat{s}}^2 \end{pmatrix}$$

$$\begin{cases} M_{\hat{h}\hat{H}}^2 = \frac{1}{2}(M_Z^2 - \lambda^2 v^2) \sin 4\beta \\ M_{\hat{H}\hat{s}}^2 = \lambda v \Lambda \cos 2\beta \\ M_{\hat{h}\hat{s}}^2 = \lambda v (2\mu - \Lambda \sin 2\beta) \end{cases}$$

Diagonal elements, $M_{\hat{h}\hat{h}}^2$, $M_{\hat{H}\hat{H}}^2$, $M_{\hat{s}\hat{s}}^2$, are more complicated.
We trade them for physical scalar masses (m_h , m_s , m_H).

- For a given $m_h \simeq 125$ GeV, m_s , m_H , μ , λ , Λ , $\tan \beta$ we can find numerically $\tilde{S}_{ij}$.

Backup slides: XENON1T – May 2017

